

<p>
</p> <p style="margin: 0cm 0cm 0pt; text-align: center; mso-layout-grid-align: none;" align="center">Ziua European? aInform?riidespre Antibiotice (ZEIA)</p> <p> </p> <p style="margin: 0cm 0cm 0pt; text-align: center; mso-layout-grid-align: none;" align="center">18 noiembrie 2018</p> <p> </p> <p style="margin: 0cm 0cm 10pt; text-align: center;" align="center"></p> <p> </p> <p style="margin: 0cm 0cm 10pt; text-align: center;" align="center">Comunicat de pres? </p> <p> Rezisten?a la antimicrobiene (RAM) continu? s? amenin?e eficien?a tratamentelor ♦potriva infec?iilor cauzate de bacterii, parazi?i, virusuri ?i fungi. Ca ilustrare a gravit?ii fenomenului, ♦ 2016, 490 000 persoane au dezvoltat la nivel global forme de tuberculoze (TB) multi-rezistente la medica?ia uzual?, iar RAM ♦cepe s? complice combaterea HIV ?i a malariei. <span lang="IT" style="color: black; line-height: 115%; font-family: "Times New Roman";" serif"; font-size: 12pt; mso-ansi-language: IT; mso-fareast-font-family:

"Times New Roman"; mso-font-ker-ning: 12.0pt; mso-fareast-language: EN-US; mso-bidi-language: AR-SA;">Costul ♦grijirii medicale pentru pacien?ii infecta?i cu microbi rezisten?i este semnificativ mai ridicat prin durata mai lung? a bolii, test?rilor suplimentare ?i nevoii de a recurge la medicamente mai scumpe. ♦ particular, rezisten?a la antibiotice (RAB) prive?te bacteriile ?i unele specii de fungi, constituind con?inutul dominant al RAM</p><p>♦</p> <p>>>> mai mult
</p>